

RU RMT 20 · PATHOLOGICAL TREATMENTS · LECTURE 19

Piriformis Syndrome

Lecture 19 — Advanced Massage Techniques for a Soft-Tissue Nerve-Compression Syndrome

How this lecture fits together



01 · Understanding

What piriformis syndrome is, its causes, and why it is so often missed.



02 · Pathophysiology

The gluteal anatomy, the sciatic nerve, and how compression creates pain.



03 · Clinical Picture

The signs, symptoms, gait changes, and what falls within our scope.



04 · Medical Management

Conservative care, injections, surgery, and medications.



05 · Massage Therapy

Assessment, goals, frequency, and a careful layer-by-layer protocol.



06 · Aftercare & Safety

Homework, contraindications, and the key cautions around the nerve.

What you will be able to do



Define the syndrome

Explain piriformis syndrome as a soft-tissue nerve-compression condition.



Trace the mechanism

Follow hypertonicity to trigger points to sciatic compression and pain.



Recognise the picture

Identify the symptoms and gait changes, and rule-out mimics like sciatica.



Assess within scope

Gather the right intake and observational findings before treating.



Treat layer by layer

Apply a gentle myofascial, static-compression protocol.



Work safely

Apply the contraindications that protect a fragile, compressed nerve.

A soft-tissue nerve-compression syndrome

Piriformis syndrome occurs when the piriformis muscle — deep in the buttock — compresses, irritates, or entraps the nearby sciatic nerve. The result is pain, numbness, and tingling that can radiate down the leg.

Because it mimics low-back pain from a disc problem, it is frequently misdiagnosed. Yet it is fundamentally a soft-tissue condition — which places much of its symptom relief squarely within our scope.



A hypertonic muscle

Piriformis spasm irritates the nerve.



Sciatic-nerve pain

Radiating buttock and leg pain.



Often misdiagnosed

Mistaken for disc-related LBP.



Within our scope

A soft-tissue condition we can help.

Source: Rattray & Ludwig (2000); Ruhani College Lecture 19



01

SECTION ONE

Understanding Piriformis Syndrome

A small deep muscle, a large fragile nerve, and a long list of imitators — understanding this syndrome begins with knowing why it is so easily missed.

What causes piriformis syndrome

 **Muscle spasm or hypertonicity.** A spasming or extremely hypertonic piriformis compresses, irritates, or injures the proximal sciatic nerve.

 **Trauma & fibrosis.** Direct injury or traumatic fibrosis of the muscle can entrap the nerve.

 **Postural load.** Hyperlordosis and anatomic abnormalities alter the muscle's pull on the nerve.

 **Prolonged sitting.** Sustained pressure through the buttock — a common daily driver.

 **Explosive movement.** Vigorous activity involving sudden bending or twisting can trigger it.

Source: Rattray & Ludwig (2000); Ruhani College Lecture 19

Why it is so often misdiagnosed



Piriformis syndrome is frequently mistaken for clinical low-back pain with radiculopathy — pain radiating from the spine — caused by a lumbar disc problem. Its symptoms overlap almost perfectly.



Several physicians. A client may see many clinicians before the correct diagnosis is reached.



Unnecessary surgery. Misdiagnosis can lead to inappropriate treatment — even needless spinal disc surgery.

Source: Rattray & Ludwig (2000)

Populations & patterns



Athletes

Skiers, tennis players, and long-distance cyclists — sports with repetitive hip loading.



More common in women

The condition shows a notable prevalence among women.



Prolonged sitters

Truck drivers and taxi drivers, from hours of sustained buttock pressure.



Onset & duration

Gradual when lifestyle-related, sudden when trauma-related; typically lasting weeks to months.

Source: Rattray & Ludwig (2000); Ruhani College Lecture 19

A leading contributor to low-back pain

30–45%

of people aged 18–55 experience some form of low-back pain

~50%

of piriformis cases report a history of direct trauma to the buttock

Piriformis syndrome ranks alongside sciatic-nerve entrapment, herniated disc, direct trauma, and muscle spasm as a leading cause of low-back pain. Reported trauma includes motor-vehicle accidents, a fall onto the buttock, difficult childbirth, or a hip or low-back torsion injury. Common comorbidities are degenerative lumbar disc disease, ischial-tuberosity bursitis, and sciatica.

Source: Rattray & Ludwig (2000); Ruhani College Lecture 19

Why accurate diagnosis matters most

The most serious complications don't come from the syndrome itself — they come from getting the diagnosis wrong. Misdiagnosis can worsen symptoms, invite inappropriate treatment, and lead to unnecessary surgery, long-term disability, and pain.

The prognosis is good when diagnosis is early and accurate, and recurrence is uncommon once a rigorous course of therapy is followed.



Improper diagnosis

The main source of serious harm.



Needless surgery

An avoidable, high-cost outcome.



Good with early care

Accurate, timely treatment helps.



Low recurrence

When rigorous therapy is followed.

Source: Rattray & Ludwig (2000)

How piriformis syndrome is identified

 **Physical assessment & history.** Diagnosis rests on examination plus the neurologic history and any pelvic trauma or childbirth difficulty.

 **Manual compression.** The muscle is accessible through the rectum; manual compression that reproduces the pain helps confirm the condition.

 **MR neurography.** A newer technique that accurately reveals piriformis asymmetry and sciatic-nerve hyperintensity.

 **Rule-out imaging.** X-ray, MRI, and nerve-conduction tests exclude other conditions with similar symptoms.

There is no single definitive test — diagnosis is a process of confirmation and exclusion.



SECTION TWO

Pathophysiology

Picture the anatomy clearly: a deep rotator muscle lying directly over a large but fragile nerve — and a pain signal born of oxygen starvation.

02

The gluteal complex & the piriformis

Clearly envisioning the anatomy of the gluteal complex is the therapist's most important tool for this syndrome. The piriformis is a deep hip rotator that lies beneath the thick gluteus maximus.

Running immediately alongside — and in some people through — the muscle is the proximal sciatic nerve: large, but surprisingly fragile under pressure.



Deep to glute max

The piriformis sits beneath thick muscle.



A hip rotator

It laterally rotates the hip.



Beside the nerve

The sciatic nerve runs alongside it.



A fragile nerve

Large, but vulnerable to compression.

Source: Rattray & Ludwig (2000); Ruhani College Lecture 19

The piriformis and the sciatic nerve



Hypertonicity

A chronically tight piriformis presses on the proximal sciatic nerve.



Nerve compression

Those tissues compress and irritate the nerve where it exits the pelvis.



Trigger points

Sustained tension seeds myofascial trigger points in and around the muscle.



Radiating pain

The nerve refers pain, numbness, and tingling down the leg.

Source: Rattray & Ludwig (2000); Travell & Simons (1999)

How tightness becomes nerve pain



Hypertonicity

The piriformis and neighbours tighten and shorten.



Trigger points

Myofascial trigger points develop in the tense tissue.



Nerve compression

The sciatic nerve is compressed and irritated.

Massage therapy works along this whole cascade — softening the tissue layer by layer to relieve the compression at its source.

Source: Rattray & Ludwig (2000)

The oxygen-deprivation pain signal



A nerve deprived of oxygen through compression or impingement alerts the body to its need for oxygen by signalling pain — often surprisingly intense pain. It is the same mechanism seen in the pain-spasm-pain cycle and in TMJ syndrome.



Compression starves the nerve. Reduced blood flow means reduced oxygen.



Pain is the alarm. The nerve's distress signal is felt as pain along its path.

Source: Rattray & Ludwig (2000)

Chronic, low-level compression

Blunt-force trauma is a common cause — but it is far from the only one. Even low-level, chronic compression, sustained over years, can produce piriformis syndrome.

The classic example is a wallet kept in the same back pocket for years, quietly pressing on the nerve every time its owner sits down.



Acute trauma

A fall or blow to the buttock.



Chronic pressure

Low-level load over years.



The wallet effect

A back-pocket wallet while sitting.



Same result

The fragile nerve is compressed.

Source: Rattray & Ludwig (2000)



03

SECTION THREE

The Clinical Picture

The symptoms range from local buttock pain to radiating numbness and subtle changes in gait — and much of the picture falls within our scope.

What the client reports



Deep regional pain. In the buttock, coccyx, hip, groin, or posterior thigh.



Radiating numbness & tingling. Travelling down the posterior leg and into the lateral foot.



Worse with sitting. Prolonged sitting, climbing stairs, or squatting all aggravate it.



Piriformis spasm. Palpable spasm and hypertonicity, with a tender gluteal region.



Weak hip abduction. Reduced strength moving the leg outward.



SI-joint hypomobility. A stiff sacroiliac joint, sometimes with pain on bowel movements.

Source: Rattray & Ludwig (2000); Ruhani College Lecture 19

Gait & postural signs



Shortened stride

The client takes a noticeably shorter step on the affected side.



Apparent leg shortening

The leg on the affected side can appear shorter.



A splayed foot

Marked lateral rotation of the foot when the client lies supine.

These quiet, observable signs — seen as the client walks, sits, and lies down — often tell you as much as the reported pain.

Source: Rattray & Ludwig (2000); Ruhani College Lecture 19

Piriformis syndrome vs. sciatica



Piriformis syndrome

A muscular entrapment

- A soft-tissue problem in the buttock
- The piriformis compresses the nerve
- Well suited to massage therapy
- Reproduced by muscle compression



Sciatica

A nerve-root problem

- Often arises from the lumbar spine
- A disc or root is the source
- May need medical management
- Anatomically related, but distinct

The two are frequently confused — they are anatomically related, but they are genuinely different conditions.

Source: Rattray & Ludwig (2000)

Symptoms massage therapy can address

Piriformis syndrome is a soft-tissue condition, so much of its symptom relief sits well within the massage therapist's scope of practice.



Relieve nerve compression

Layer-by-layer softening of hypertonic muscle unloads the nerve.



Ease pain & tingling

Multiple techniques address symptoms of nerve oxygen deprivation.



Reduce muscle tenderness

Careful muscle-stimulating techniques calm the tender tissue.



Lower anxiety

Relaxing work helps the client reach a parasympathetic state and improves quality of life.

Source: Rattray & Ludwig (2000); Ruhani College Lecture 19



04

SECTION FOUR

Medical Management

Care begins conservatively and escalates only when it must — and the massage therapist works as one member of a physician-led team.

From conservative care to surgery



Conservative first. Spinal/pelvic traction, progressive stretching and strengthening, careful ROM, heat and cold, ultrasound, and activity modification.



Daily home self-care. A rigorous, daily self-care regimen is essential for complete recovery.



Corticosteroid injection. Injected into the muscle belly only when conservative methods fail.



Surgery, last of all. Reserved as the treatment of last resort.

A sports-medicine or orthopedic physician most often leads treatment — frequently after the client has seen several clinicians.

Managing pain & inflammation



NSAIDs

Non-steroidal anti-inflammatories such as ibuprofen (Motrin, Advil) reduce pain and inflammation.



A treatment caution

Avoid cross-fibre friction while a client is taking anti-inflammatory medication.



Local anaesthetics

Agents such as bupivacaine and topical lidocaine (Lidoderm, Xylocaine) numb the painful region.



Pain can be masked

When medication dulls pain, moderate your pressure and rely on frequent feedback.

Source: Rattray & Ludwig (2000); Werner (2019)

The massage therapist's place in care

The massage therapist does not diagnose piriformis syndrome. That diagnosis belongs to the physician; our role is to treat the secondary soft-tissue symptoms it produces.

That means working closely with the sports-medicine or orthopedic physician and the physical therapist who already know the exact anatomic picture.



Physician-diagnosed

We treat, we do not diagnose.



Alongside the PT

Coordinate with the care team.



Treat the symptoms


Focus on the soft-tissue effects.



Stay in scope

Respect the boundaries of practice.

Source: Rattray & Ludwig (2000); Ruhani College Lecture 19



05

SECTION FIVE

Massage Therapy Treatment

Assess carefully, then work slowly — warming and softening each layer with static compression until the piriformis itself can be reached and released.

Questions to ask before you begin



Who is on the care team? Is the client under a sports-medicine, orthopedic, or chiropractic physician?



Any recent injection? Has she had a corticosteroid or anaesthetic injection into the buttock in the last 10 days?



Any recent rectal procedure? Has she had a rectal examination or treatment in the last 10 days?



What is the pain now? Ask her to rate her current pain on a scale of 0 to 10.

Source: Rattray & Ludwig (2000); Ruhani College Lecture 19

Reading the pain & the movement



The path of the pain. Trace exactly where the pain travels.



The nature of the pain. Is it burning, numb, tingling, dull, or achy?



Guarding as she walks. Watch for protective guarding, an apparent limp, or signs of leg shortening.



Bracing at the chair. Note whether she braces herself when sitting down or rising.

Source: Rattray & Ludwig (2000); Ruhani College Lecture 19

What we aim to achieve

Because piriformis syndrome is a soft-tissue condition, the therapist — working closely with the physician and physical therapist — has clear, achievable goals.



Unload the nerve

Reduce the compressive force on the affected sciatic nerve.



Improve posture

Facilitate better posture to ease ongoing strain.



Reduce hypertonicity

Soften the piriformis and the surrounding gluteal muscles.



Provide pain relief

Deliver genuine, lasting relief of the client's pain.

Source: Rattray & Ludwig (2000); Ruhani College Lecture 19

A general treatment schedule

2× / wk

60-minute sessions in the early, painful stages

1× / wk

60-minute sessions as the symptoms begin to subside

Monthly

60-minute maintenance until symptoms fully resolve

Consistency, paired with the client's daily home regimen, is what carries the recovery through to completion.

Source: Rattray & Ludwig (2000); Ruhani College Lecture 19

The guiding philosophy

Effective treatment depends on clearly visualising the structures beneath your hands, and a willingness to begin gently and move, layer by layer, only to the depth the client can tolerate.

Although deep trigger points are common, the protocol does not call for aggressive trigger-point work. It leans on myofascial technique — working muscle and fascia — and warming superficial tissue first.



See the structures

Visualise the layers you work.



Layer by layer

To the depth she tolerates.



Gentle, not aggressive

No forceful trigger-point work.



Static compression

Broad open hand and forearm.

Source: Rattray & Ludwig (2000); Ruhani College Lecture 19

Static compression, done safely



Light to deep, then wait. Apply direct, non-moving pressure, progressing gently and waiting for the tissue to release, soften, or shift ever so slightly.



Let the tissue lead. That small movement is the body's acquiescence — your signal that you may go deeper.



No elbows, knuckles, or fists. An untrained elbow with poor mechanics can damage the already-fragile sciatic nerve.



Work toward the origin. Know the muscles' origins and insertions, and work toward the origin to soften and prevent spasm.

Source: Rattray & Ludwig (2000); Ruhani College Lecture 19

Locating the piriformis



With the client side-lying, palpate the posterior-superior iliac spine (PSIS) and the greater trochanter. Lay the heel of your hand on the PSIS and point your fingertips toward the greater trochanter.



Follow the line. The piriformis lies deep to the gluteals along that PSIS-to-trochanter pathway.



Deep to glute max. You must work through the thick gluteus maximus to reach it.

Source: Rattray & Ludwig (2000); Ruhani College Lecture 19

Positioning, draping & hydrotherapy



Hydrotherapy ready

Hot packs for dull, aching or distal tissue; cold packs to quiet a spasm or sharp, stabbing pain.



Draping is paramount

Work involves the hip and gluteal complex; skin-on-skin is most effective, with careful, respectful draping.



Comfortable side-lying

Provide plenty of pillows to support a relaxed side-lying position.



Help her relax

A client in pain guards herself; a calm environment and a few deep breaths help her settle.

Source: Rattray & Ludwig (2000); Ruhani College Lecture 19



06

SECTION SIX

Aftercare & Safety

Here, less is more: keep self-care simple, defer to the physical therapist's regimen, and respect the firm cautions that protect the nerve.

Simple, safe self-care

Unlike most conditions, here you deliberately keep self-care simple — the risk of harm means you offer lifestyle suggestions and encourage the rigorous regimen the client's PT has already assigned.



Sit smarter

For long sitting, use a padded or rocking chair; when driving, get out and walk every hour.



Pillow support

A pillow between the knees when side-lying, and under the knees when supine, offloads the low back.



Ice or heat

Ice for acute, sharp local pain; heat for dull, aching pain.



Gentle movement

Yoga or tai chi can help — but only with the physician's and PT's permission.

Source: Rattray & Ludwig (2000); Ruhani College Lecture 19

Protecting a compressed nerve



Modulate elbow work. The sciatic and neighbouring nerves are already compressed; an ill-placed, aggressive elbow can cause serious damage.



No cross-fibre friction on anti-inflammatories. Skip cross-fibre friction while the client is taking anti-inflammatory medication.



No active or passive joint ROM. The PT and physician know the exact anatomic damage; without it, even innocuous ROM can harm.



No hip ROM in pregnancy with unexplained LBP. Especially avoid it in the third trimester when the cause of the pain is unknown.

Source: Rattray & Ludwig (2000); Werner (2019)

Sources & further reading

- Rattray, F., & Ludwig, L. (2000). Clinical Massage Therapy: Understanding, Assessing and Treating Over 70 Conditions. Toronto: Talus Inc.
- Travell, J. G., & Simons, D. G. (1999). Myofascial Pain and Dysfunction: The Trigger Point Manual (2nd ed.). Baltimore: Williams & Wilkins.
- Werner, R. (2019). A Massage Therapist's Guide to Pathology (7th ed.). Books of Discovery.
- Magee, D. J. (2014). Orthopedic Physical Assessment (6th ed.). St. Louis: Saunders / Elsevier.
- Lowe, W. (2009). Orthopedic Assessment in Massage Therapy. Sisters, OR: Daviau-Scott.
- Ruhani College. Lecture 19 — Piriformis Syndrome (course notes).



KEY TAKEAWAYS

Work slow, work safe, respect the nerve

Piriformis syndrome is a soft-tissue nerve-compression condition that hides behind low-back pain. Diagnosed by a physician and treated by our hands, it responds to patient, layer-by-layer myofascial work — gentle static compression, no aggressive elbows, and constant client feedback to protect the fragile sciatic nerve.

Ruhani College · RU-RMT 20 Pathological Treatments · Lecture 19 — Piriformis Syndrome